

CONVERSE THEOREMS AND EXTENSIONS  
IN CHEBYSHEV RATIONAL APPROXIMA-  
TION TO CERTAIN ENTIRE FUNC-  
TIONS IN  $[0, +\infty)$

BY

G. MEINARDUS, A. R. REDDY, G. D. TAYLOR AND  
R. S. VARGA

Reprinted from the  
BULLETIN OF THE AMERICAN MATHEMATICAL SOCIETY  
May, 1971, Vol. 77, No. 3  
Pp. 460-461

CONVERSE THEOREMS AND EXTENSIONS IN  
CHEBYSHEV RATIONAL APPROXIMATION TO  
CERTAIN ENTIRE FUNCTIONS IN  $[0, +\infty)$

BY G. MEINARDUS,<sup>1</sup> A. R. REDDY,<sup>2</sup> G. D. TAYLOR<sup>3</sup> AND R. S. VARGA<sup>4</sup>

Communicated by Eugene Isaacson on November 19, 1970

Recently, it has been shown that the problem of rational approximation to  $e^{-z}$  in  $[0, +\infty)$  arises naturally in numerical methods for approximating solutions of heat-conduction-type partial differential equations [1], [2]. This special approximation problem leads one to the general question of approximating functions on the half line  $[0, +\infty)$ . In this paper we wish to announce two results of this study which are in the spirit of work done by S. N. Bernstein. A complete description of this work with proofs of these results and additional results will appear elsewhere.

In order to state these results, we need the following notation. For any nonnegative integer  $m$ , let  $\pi_m$  denote the collection of all real polynomials of degree at most  $m$ . For given  $r > 0$  and  $s > 1$ , let  $\mathcal{E}(r, s)$  denote the unique open ellipse in the complex plane with foci at  $x=0$  and  $x=r$  and semimajor and semiminor axes  $a$  and  $b$  such that  $b/a = (s^2 - 1)/(s^2 + 1)$ . Finally, if  $f(z)$  is any entire function, we set

$$\tilde{M}_f(r, s) = \sup\{|f(z)| : z \in \mathcal{E}(r, s)\}$$

and

$$M_f(r) = \sup\{|f(z)| : |z| = r\}.$$

We now state our main results.

**THEOREM 1.** *Let  $f(x)$  be a real continuous function on  $[0, +\infty)$  with at most a finite number of points  $\{x_i\}_{i=1}^m$  in  $[0, +\infty)$  for which  $f(x_i) = 0$ . Assume that there exist a sequence of real polynomials  $\{p_n(x)\}_{n=0}^{\infty}$ , with  $p_n \in \pi_n$  for each  $n \geq 0$ , and a real number  $q > 1$  such that*

*AMS 1970 subject classifications.* Primary 41A20; Secondary 41A50, 41A25.

*Key words and phrases.* Approximation theory, rational approximation, approximation of entire functions, geometric convergence.

<sup>1</sup> Present address—University of Erlangen-Nurnberg, Erlangen, Germany.

<sup>2</sup> Present address—University of Toledo, Toledo, Ohio 43606.

<sup>3</sup> Supported in part by NSF Grant CP-12088.

<sup>4</sup> Supported in part by AEC Grant AT (11-1)-2075; present address—Kent State University, Kent, Ohio 44240.

$$\limsup_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{1}{p_n(x)} \right\|_{L_\infty[0, \infty)} \right\}^{1/n} = \frac{1}{q} < 1.$$

Then, there exists an entire function  $F(z)$  with  $F(x) = f(x)$  for all  $x \geq 0$ , and  $F$  is of finite order  $\rho$ , i.e.,

$$\limsup_{r \rightarrow \infty} \frac{\ln \ln M_F(r)}{\ln r} \equiv \rho < \infty.$$

In addition, for each  $s > 1$ , there exist real numbers  $K = K(q, s) > 0$ ,  $\theta = \theta(q, s) > 1$  and  $r_0 = r_0(q, s) > 0$  such that  $\bar{M}_F(r, s) \leq K(\|f\|_{L_\infty[0, r]})^\theta$  for all  $r \geq r_0$ . If, for each  $s > 1$ ,  $\bar{\theta}(s)$  is defined by

$$\limsup_{r \rightarrow \infty} \left\{ \frac{\ln \bar{M}_F(r, s)}{\ln \|f\|_{L_\infty[0, r]}} \right\} = \bar{\theta}(s)$$

when  $\|f\|_{L_\infty[0, r]}$  is unbounded as  $r \rightarrow \infty$ , and  $\bar{\theta}(s) \equiv 1$  otherwise, then the order  $\rho$  of  $F$  satisfies

$$\rho \leq \inf_{s > 1} \left\{ \frac{\ln \bar{\theta}(s)}{\ln \left[ \frac{1}{2} + \frac{1}{4}(s + 1/s) \right]} \right\},$$

and this upper bound for the order  $\rho$  is in general best possible.

**THEOREM 2.** Let  $f(z) = \sum_{k=0}^{\infty} a_k z^k$  be an entire function with non-negative Taylor coefficients and  $a_0 > 0$ . If there exist real numbers  $s > 1$ ,  $A > 0$ ,  $\theta > 0$  and  $r_0 > 0$  such that  $\bar{M}_f(r, s) \leq A(\|f\|_{L_\infty[0, r]})^\theta$  for all  $r \geq r_0$  then there exist a sequence of real polynomials  $\{p_n(x)\}_{n=0}^{\infty}$  with  $p_n \in \pi_n$  for each  $n \geq 0$ , and a real number  $q \geq s^{1/(1+\theta)} > 1$  such that

$$\limsup_{n \rightarrow \infty} \left\{ \left\| \frac{1}{f(x)} - \frac{1}{p_n(x)} \right\|_{L_\infty[0, \infty)} \right\}^{1/n} = \frac{1}{q} < 1.$$

The requirement that the Taylor coefficients be nonnegative in Theorem 2 is due to a technical difficulty in the proof of the theorem. Examples can be given to show that this requirement is not necessary for geometric convergence to occur.

#### REFERENCES

1. W. J. Cody, G. Meinardus and R. S. Varga, *Chebyshev rational approximations to  $e^{-x}$  in  $[0, +\infty)$  and applications to heat-conduction problems*, J. Approximation Theory 2 (1969), 50-65. MR 39 #6536.
2. G. Meinardus and R. S. Varga, *Chebyshev rational approximation to certain entire functions in  $[0, +\infty)$* , J. Approximation Theory 3 (1970), 300-309.

MICHIGAN STATE UNIVERSITY, EAST LANSING, MICHIGAN 48823