

Circle one: 9:55 / 12:05

Dr. Kracht

Print Name: KEY

Exam 3 Version A
Friday, October 24, 2014

NO CALCULATORS.

Academic Honesty Pledge

Your signature at the bottom indicates your agreement to abide by the following rules.

1. I have placed all purses, bags, books, notes, and other papers in the designated area of the classroom.
2. **I have placed all cell phones, calculators, and other electronic devices in the designated area of the classroom.**
3. I will not communicate with other students during the exam.
4. I will not seek help from or give help to others during the exam.
5. I will turn my exam in and will not take it from the classroom.
6. I will not discuss the exam outside of class with another student who has not yet taken the exam.
7. I will not cheat in any other way.
8. I will follow any other instructions from my professor.

Signature: _____

Good Luck!

Print Name: KEY Exam Score: /100
(105 pts available)

Exam 3: Version A
NO CALCULATORS.

Part I: Long Answer.

No credit for answers without sufficient justification. Use standard mathematical notation correctly.

1. (20 points) Evaluate each of the following by hand. Write each answer as a whole number or fraction in lowest terms. No credit for answers without all steps written out clearly.

$$3 \quad (a) \frac{569!}{568!} = \frac{569 \cdot (568!)}{568!}$$

$$= \underline{569}.$$

$$3 \quad (b) \frac{6!}{6} = \frac{6 \cdot 5!}{6}$$

$$= 5!$$

$$= 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1$$

$$= \underline{120}.$$

$$4 \quad (c) \frac{10!}{4! 6!} = \frac{10 \cdot 9 \cdot 8 \cdot 7 \cdot (6!)}{(4 \cdot 3 \cdot 2 \cdot 1) (6!)}$$

$$= \frac{10 \cdot \overset{3}{\cancel{9}} \cdot \cancel{8} \cdot 7}{4 \cdot \cancel{3} \cdot \cancel{2}}$$

$$= 10 \cdot 3 \cdot 7$$

$$= 10 \cdot 21$$

$$= \underline{210}.$$

$$5 \quad (d) {}_7C_2 = \frac{7!}{(7-2)! 2!}$$

$$= \frac{7!}{5! 2!}$$

$$= \frac{7 \cdot 6 \cdot 5!}{5! (2 \cdot 1)}$$

$$= \frac{7 \cdot 6}{2}$$

$$= 7 \cdot 3$$

$$= \underline{21}.$$

$$5 \quad (e) {}_7P_2 = \frac{7!}{(7-2)!}$$

$$= \frac{7 \cdot 6 \cdot (5!)}{5!}$$

$$= 7 \cdot 6$$

$$= \underline{42}.$$

2. (25 points) Consider an six-sided die (with sides labeled "1," "2," ..., "6") and a four-sided die (with sides labeled "1," "2," "3," and "4"). The dice are tossed and we observe the number that comes up on each die.

- (a) Write out the sample space S for this random experiment completely. Record each outcome as an ordered pair of the form (s, f) where s is the number that comes up on the six-sided die and f is the number that comes up on the four-sided die.

$$S = \{ (1,1), (1,2), (1,3), (1,4), \\ (2,1), (2,2), (2,3), (2,4), \\ (3,1), (3,2), (3,3), (3,4), \\ (4,1), (4,2), (4,3), (4,4), \\ (5,1), (5,2), (5,3), (5,4), \\ (6,1), (6,2), (6,3), (6,4) \}$$

- (b) Let E_1 be the event "roll doubles or a total of four."

- i. Write E_1 as a set.

$$E_1 = \{ (1,1), (2,2), (3,3), (4,4), (1,3), (3,1) \}$$

- ii. Find $|E_1|$. = 6

- iii. Find $\Pr(E_1)$. = $\frac{6}{24} = \frac{1}{4}$

- (c) Let E_2 be the event "roll a total of twelve."

- i. Write E_2 as a set.

$$E_2 = \{ \}$$

- ii. Find $|E_2|$. = 0

- iii. Find $\Pr(E_2)$. = $\frac{0}{24} = 0$.

Part II: Multiple Choice (5 points each)

Circle the letter of the best answer.

E

3. Which of the following is not a random experiment?

- (a) rolling a pair of dice
- (b) drawing a card from a deck of playing cards
- (c) tossing a coin three times
- (d) selecting numbers in a lottery
- (e) All of these are random experiments

D

4. In the game of Bunco, three ordinary six-sided dice are rolled and the number on each die is observed. What is the size of the sample space?

- $6 \cdot 6 \cdot 6 = 6^3$
- (a) 3^6
 - (b) $6!$
 - (c) 6C_3
 - (d) 6^3
 - (e) none of these

C

5. A person shoots 4 consecutive free throws and the total number of successes is observed. What is the size of the sample space?

- $S = \{0, 1, 2, 3, 4\}$
 $|S| = 5$
- (a) 2^4
 - (b) $4!$
 - (c) 5
 - (d) 4^2
 - (e) none of these

A

6. Johnie's Coffee Shop offers platters consisting of a sandwich, salad, vegetable, and drink. There are eight different sandwiches, three salads, five vegetables, and ten drinks to choose from. How many different platters does Johnie's offer?

- Multiplication Rule:**
 $8 \cdot 3 \cdot 5 \cdot 10 = 24 \cdot 5 \cdot 10$
 $= 120 \cdot 10$
 $= 1200$
- (a) 1200
 - (b) 26
 - (c) 130
 - (d) 2000
 - (e) none of these

7. Eight children are trick-or-treating together. In how many ways can they line up at the door to receive their treats?

B

- (a) 2^8
- (b) $8!$
- (c) ${}_8 C_8$
- (d) 8
- (e) none of these

$$\underline{8} \cdot \underline{7} \cdot \underline{6} \cdot \underline{5} \cdot \underline{4} \cdot \underline{3} \cdot \underline{2} \cdot \underline{1} = 8!$$

8. Twenty contestants are entered in a Halloween costume contest. In how many ways can one choose the top four finishers regardless of order?

B

- (a) ${}_{20} P_4$
- (b) ${}_{20} C_4$
- (c) $4!$
- (d) 20^4
- (e) none of these

↳ combination

9. Twenty contestants are entered in a Halloween costume contest. In how many ways can one choose Best Costume and the first-, second-, and third-runners-up?

A

- (a) ${}_{20} P_4$
- (b) ${}_{20} C_4$
- (c) $4!$
- (d) 20^4
- (e) none of these

↳ permutation

10. In a probability space, which of the following statements is not necessarily true?

A

- (a) All outcomes are equally likely.
- (b) The probability of the sample space is equal to 1.
- (c) The probability of an outcome is always between 0 and 1 (inclusive).
- (d) The probability of the impossible event is 0.
- (e) All of these statements are always true.

Version A

11. Consider the sample space $S = \{o_1, o_2, o_3, o_4\}$. Suppose $\Pr(o_1) = 0.30$ and $\Pr(o_2) = 0.50$. If o_4 is four times as likely as o_3 , find $\Pr(o_4)$.

C

- (a) 0.10
- (b) 0.40
- (c) 0.16
- (d) 0.20
- (e) none of these

Given: $\Pr(o_4) = 4 \Pr(o_3)$.

Let $\Pr(o_3) = p$. Then $\Pr(o_4) = 4p$.

We always have:

$$\Pr(o_1) + \Pr(o_2) + \Pr(o_3) + \Pr(o_4) = 1$$

So $0.30 + 0.50 + p + 4p = 1$

$$5p = 1 - 0.80$$

$$5p = 0.20$$

$$p = 0.04$$

So $\Pr(o_4) = 4p = 0.16$.

12. An honest coin is tossed five times. What is the probability of tossing 2 heads and 3 tails? event E

A

- (a) $\frac{{}^5C_2}{2^5}$
- (b) $\frac{{}^5P_2}{2^5}$
- (c) $\frac{2}{2^5}$
- (d) $\frac{{}^5C_2}{5!}$
- (e) none of these

$$|S| = 2^5$$

$|E| = {}^5C_2$: # of ways of selecting the two slots for the heads

So $\Pr(E) = \frac{|E|}{|S|} = \frac{{}^5C_2}{2^5}$

13. In how many ways can one distribute 3 ghost costumes, 5 vampire costumes, and 2 witch costumes among ten children?

B

- (a) $({}_{10}C_3) \cdot ({}_{10}C_5) \cdot ({}_{10}C_2)$
- (b) $({}_{10}C_3) \cdot ({}_{7}C_5) \cdot ({}_{2}C_2)$
- (c) $({}_{10}P_3) \cdot ({}_{7}P_5) \cdot ({}_{2}P_2)$
- (d) $3! + 5! + 2!$
- (e) none of these

${}_{10}C_3$ ways to choose the ghosts

(so 7 children left)

${}_{7}C_5$ ways to choose the vampires

(then 2 children left)

${}_{2}C_2 = 1$ way to choose the witches.

Then use Multiplication Rule.

14. How many permutations are there of the letters in the word "HALLOWEEN"?

D

- (a) $9!$
- (b) $({}_9P_2) \cdot ({}_7P_2) \cdot ({}_5P_5)$
- (c) $({}_9C_2) \cdot ({}_7C_2) \cdot ({}_5C_5)$
- (d) $({}_9C_2) \cdot ({}_7C_2) \cdot (5!)$
- (e) none of these

${}_9C_2$ ways to place the 2 L's;

Then

${}_7C_2$ ways to place the 2 E's;

Then ${}_5P_5 = 5!$ ways to order the remaining 5 letters.