

Name: _____ Score: _____ /100
(113 pts available)

EXAM 3: Version A

NO CALCULATORS.

Multiple Choice: 17 questions at 4 points each.

Circle the letter of the best response.

Although your work will not be graded, you should write out complete and careful solutions to reduce the chance of error.

1. The lowest point of the graph of $f(x) = x^3 - 3x + 1$ on the interval $[-1, 3]$ is

- (a) $(2, -9)$
- (b) $(0, -3)$
- (c) $(1, -1)$
- (d) $(-1, 0)$
- (e) none of these

2. The highest point of the graph of $f(x) = x^3 - 3x + 1$ on the interval $[-1, 3]$ is

- (a) $(3, 19)$
- (b) $(0, 1)$
- (c) $(-1, 3)$
- (d) $(3, 24)$
- (e) none of these

3. True or False: Suppose f is continuous and that $f'(7) = 0$ and $f''(7) = -10$. Then we may conclude that f has an **absolute maximum** point at $x = 7$.

- (a) True
- (b) False

4. True or False: If f is continuous on the interval $[1, 5]$, then f must have an **absolute minimum value** on $[1, 5]$.
- (a) True
 - (b) False
-

5. True or False: If f is continuous on the interval $(1, 5)$, then f must have an **absolute minimum value** on $(1, 5)$.
- (a) True
 - (b) False
-

6. An automobile dealer can sell 8 cars per day at a price of \$25,000. He estimates that for each \$1000 price reduction, he can sell 3 more cars each day. Let x be the number of \$1000 price reductions. Find an expression for the price per car (in dollars).
- (a) $25,000x - 1000$
 - (b) $25,000 \times 1000x$
 - (c) $25,000 - 8x$
 - (d) $25,000 - 1000x$
 - (e) none of these
-

7. An automobile dealer can sell 8 cars per day at a price of \$25,000. He estimates that for each \$1000 price reduction, he can sell 3 more cars each day. Let x be the number of \$1000 price reductions. Find an expression for the number of cars sold per day.
- (a) $8 + 3x$
 - (b) $8 - 1000x$
 - (c) $8x$
 - (d) $8 - 3x$
 - (e) none of these
-

8. If the reproduction function for Albacore tuna is $f(p) = 5\sqrt[3]{p^2}$ (where both p and $f(p)$ are in thousands), find the sustainable yield for a population of 8000 tuna. (Evaluate $Y(8)$ where $Y(p)$ is the sustainable yield function.)
- (a) 8,000 tuna
 - (b) 12,000 tuna
 - (c) 20,000 tuna
 - (d) 16,000 tuna
 - (e) none of these
-

9. Evaluate the integral: $\int 2.4x \, dx$

- (a) $2.4x^2 + C$
- (b) $4.8x^2 + C$
- (c) $1.2x^2 + C$
- (d) $2.4 + C$
- (e) none of these

10. Evaluate the integral: $\int \frac{5}{x^4} dx$

- (a) $\frac{1}{x^5} + C$
 - (b) $-\frac{15}{x^3} + C$
 - (c) $-\frac{5}{3x^3} + C$
 - (d) $5 \ln |x^4| + C$
 - (e) none of these
-

11. Evaluate the integral: $\int 8e^{2t} dt$

- (a) $16e^{2t} + C$
 - (b) $4e^{2t} + C$
 - (c) $16e^{3t} + C$
 - (d) $8e^{2t} + C$
 - (e) none of these
-

12. Evaluate the integral: $\int \frac{6x^2 - 4x + 7}{x^2} dx$

- (a) $6x - 4 \ln |x| - \frac{7}{x} + C$
 - (b) $6x - 4 \ln |x| - \frac{7}{3x^3} + C$
 - (c) $6x - 4 \ln |x| - \frac{21}{x^3} + C$
 - (d) $\frac{2x^3 - 2x^2 + 7x}{\frac{1}{3}x^3} + C$
 - (e) none of these
-

13. Black squirrel flu is spreading on campus at a rate of $10e^{0.5t}$ new cases per day, where t is the number of days since the start of the epidemic. The epidemic began with 1 case. Find an expression for the total number of flu cases during the first t days.

- (a) $20e^{0.5t} + 1$
- (b) $20e^{0.5t} - 19$
- (c) $10e^{0.5t} + 1$
- (d) $5e^{0.5t}$
- (e) none of these

14. Evaluate the integral: $\int_e^{e^4} \frac{10}{x} dx$.

- (a) 0
 - (b) 3
 - (c) 30
 - (d) e^3
 - (e) none of these
-

15. Find the area under the curve $y = x^2 + 4x$ from $x = 0$ to $x = 2$.

- (a) 12
 - (b) $\frac{20}{3}$
 - (c) $\frac{32}{3}$
 - (d) $\frac{44}{3}$
 - (e) none of these
-

16. A company is considering a new manufacturing process. The rate of savings from this process is expected to be $\frac{12,000}{t^2}$ dollars per year, where t is in years. Determine the total savings from the third through the sixth year.

- (a) \$4000
 - (b) \$8000
 - (c) \$12,000
 - (d) \$2000
 - (e) none of these
-

17. Complete the statement of The Fundamental Theorem of Calculus.

Let f be continuous on $[a, b]$. Then $\int_a^b f(x) dx =$

- (a) $f(b) - f(a)$
- (b) $f'(b) - f'(a)$
- (c) $F(b) - F(a)$, where F is any antiderivative of f .
- (d) $F(a) - F(b)$, where F is any antiderivative of f .
- (e) none of these

19. (20 points) Evaluate each integral, simplifying your answer and writing out all work, including substitutions, if any. Be sure to use correct notation.

(a) $\int \sqrt{x^5 - 2} x^4 dx$

(b) $\int_{-2}^3 \frac{1}{4-x} dx$