

# Real Analysis.

Instructor: Dmitry Ryabogin

## Assignment VII.

### 1. Problem 1.

a) Let  $f(x)$  be a **monotonic** function satisfying

$$\forall x, y \in \mathbf{R}, \quad f(x) + f(y) = f(x + y), \quad f(1) = 1.$$

Prove that  $f(x) = x$ .

**Hint:** Prove that  $f(p/q) = p/q$ , then consider two **monotonic** sequences  $(r_k)_{k=1}^{\infty}$ ,  $(m_k)_{k=1}^{\infty}$ ,  $r_k, m_k \in \mathbf{Q}$ , converging to  $x \in \mathbf{R}$ ,  $x > 0$ . For  $x < 0$ , observe that  $f(x) = -f(-x)$ .

b)\* Change the word **monotonic** by **measurable** in a), and answer the same question.

**Hint:** Observe that

$$\int_0^1 f(x+y)dy = \int_0^1 (f(x) + f(y))dy = f(x) + \int_0^1 f(y)dy,$$

and conclude that  $f$  is continuous.

c)\* Is it possible to drop the measurability assumption in b)?

2. **Problem 2.** a) Let  $f(x), g(x)$  be increasing functions on the real line. Does it follow that  $f(x)g(x)$  is increasing?

b) Construct the monotonic function on the real line which is not continuous **only** in rational points.

**Hint:** Consider

$$f(x) := \sum_{k \in \mathbf{N}: r_k < x} \frac{1}{2^k}, \quad r_k \in \mathbf{Q}.$$

To show that  $f(x)$  is not continuous in rational points observe that  $\forall x > r \in \mathbf{Q}$  we have

$$f(x) = \sum_{r_k < x} \frac{1}{2^k} = \sum_{r_k < r} \frac{1}{2^k} + \sum_{r \leq r_k < x} \frac{1}{2^k}, \quad \sum_{r \leq r_k < x} \frac{1}{2^k} > \frac{1}{2^n},$$

provided  $r$  has  $n$ -th place in enumeration of rationals.

To show that  $f$  is continuous in  $\mathbf{R} \setminus \mathbf{Q}$ , observe that

$$\sup_{\mathbf{R}} f(x) = 1, \quad \inf_{\mathbf{R}} f(x) = 0,$$

hence the sum of jumps may not be bigger than 1.

*diff. for class*  
*better*

3. **Problem 3.** a) Let  $E \subset [0, 1]$ , and let  $f(x)$  be a **bounded** function on  $E$ , satisfying  $f(x_1) \leq f(x_2) \forall x_1, x_2 \in E, x_1 < x_2$ . Is it possible to extend this function in such a way that the extension  $\phi(x)$  would be nondecreasing on the whole segment  $[0, 1]$ ?

**Hint:** Let  $x_0 = \inf E$ . Put  $\phi(x) := \sup_{y < x} f(y)$  for  $x \in [0, 1] \setminus E, x > x_0$ , and  $\phi(x) := \inf_E f(y)$  for  $x \in [0, 1] \setminus E, x \leq x_0$ .

- b) Change the word **bounded** by **unbounded** in a), and answer the same question.

**Hint:** Look at "ends".

4. **Problem 4.** a) Let  $E \subset [0, 1]$  be **nowhere dense** closed set of a positive measure. Construct an **increasing** continuously differentiable  $f(x)$  on  $[0, 1]$  such that  $f'(x) = 0 \forall x \in E$ .

**Hint:** Take

$$f(x) := \int_0^x \phi(t) dt, \quad \phi(t) := \inf_{y \in E} |y - t|.$$

Then observe that  $\phi$  is continuous, and  $f'(x) = 0 \forall x \in E$ . To show that  $f$  is increasing take  $0 \leq x_1 < x_2 \leq 1$ , and observe that there exists an interval  $(\alpha, \beta) \subset (x_1, x_2)$ , such that  $(\alpha, \beta) \cap E = \emptyset$ .

- b) Is it possible to drop the assumption **nowhere dense** in a)?

**Hint:** There exists  $(\alpha, \beta) \subset E$  such that  $f'(x) = 0 \forall x \in (\alpha, \beta)$ .

5. **Problem 5.** Prove that  $\forall E \subset [0, 1], |E| = 0$ , there exists a continuous increasing function  $f(x)$  on  $[0, 1]$  such that  $\forall x \in E$  we have  $f'(x) = +\infty$ .

**Hint:**

- a)  $\forall n \in \mathbb{N}$  there exists an open bounded set  $G_n$  such that  $E \subset G_n$ , and  $m(G_n) < 2^{-n}$ .

- b) Put  $f_n(x) := \tilde{m}(G_n \cap [0, x])$ . Then  $f_n$  is increasing, continuous, nonnegative, and satisfying  $f_n(x) < 2^{-n}$ .


- c) A function  $f(x) := \sum_{n=1}^{\infty} f_n(x)$  is also increasing, continuous, nonnegative.

- d) Let  $x_0 \in E$ . Then  $[x_0, x_0 + h] \subset G_n$  for some  $n$  fixed, provided  $|h|$  is small. Prove that  $f_n(x_0 + h) = f_n(x_0) + h$ .

- e) Use d) to conclude that  $\forall k \in \mathbb{N}$  and  $|h|$  small we have

$$\frac{f(x_0 + h) - f(x_0)}{h} \geq \sum_{n=1}^k \frac{f_n(x_0 + h) - f_n(x_0)}{h} = k.$$

$f(x) = x^{1/3}$



$f(x) = \prod_{k=1}^{\infty} (x - x_k)^{1/3}$

for all  $x \in E$

for constant

leb measure  $\mathbb{R}$