

(40)

Lec 14, 2/23/09

EXAMPLES:

- ① Recall that $\langle \vec{0}, \vec{v} \rangle = 0$ for all \vec{v} in an inner product space V . It follows that $\{\vec{0}\}^\perp = V$.
- ② If V is any inner product space and $\vec{u} \in V$, then $\langle \vec{u}, \vec{v} \rangle = 0$ for all $\vec{v} \in V$. In particular, $\langle \vec{u}, \vec{u} \rangle = 0$, so by $[I_3]$, $\vec{u} = \vec{0}$. Hence $V^\perp = \{\vec{0}\}$.
- ③ Let $V = \mathbb{R}^3$ with the standard inner product, and let $S = \{(1, 2, -1), (2, -1, 2)\}$. Then

$$S^\perp = \{(x, y, z) \in \mathbb{R}^3 \mid (x, y, z) \cdot (1, 2, -1) = 0, (x, y, z) \cdot (2, -1, 2) = 0\}$$

$$= \{(x, y, z) \in \mathbb{R}^3 \mid x + 2y - z = 0, 2x - y + 2z = 0\}$$

$$= \mathcal{N}(A), \text{ where}$$

$$A = \begin{bmatrix} 1 & 2 & -1 \\ 2 & -1 & 2 \end{bmatrix} \rightarrow \begin{bmatrix} 1 & 2 & -1 \\ 0 & -5 & 4 \end{bmatrix} = E \quad \begin{array}{l} x + 2y - z = 0 \\ -5y + 4z = 0 \end{array}$$

Thus z is free, $y = \frac{4}{5}z$, $x = z - 2y = z - \frac{8}{5}z = -\frac{3}{5}z$.

Hence

$$S^\perp = \left\{ \left(-\frac{3}{5}z, \frac{4}{5}z, z \right) \mid z \in \mathbb{R} \right\} = \{ (-3r, 4r, 5r) \mid r \in \mathbb{R} \},$$

and $\{(-3, 4, 5)\}$ is a basis for S^\perp , $\dim S^\perp = 1$.

- ④ Let $V = \mathbb{R}^n$ with the standard inner product, and let A be a fixed $m \times n$ matrix, so $\mathcal{N}(A) \subseteq \mathbb{R}^n$. Then

$$x \in \mathcal{N}(A) \Leftrightarrow Ax = \vec{0}$$

\Leftrightarrow The dot product of each row of A with \vec{x} is 0

$\Leftrightarrow \vec{x}$ is orthogonal to each row of A

$\Leftrightarrow \vec{x} \in \mathcal{R}(A)^\perp$

Hence (ignoring the distinction between row vectors in $\mathcal{R}(A)$ and column vectors in $\mathcal{N}(A)$) we have

$$\mathcal{R}(A)^\perp = \mathcal{N}(A).$$

Exercises: Show that the left null space of A is the orthogonal complement of the column space of A .