

## Functions of Real Variables II (62052/72052)

Home Work 6, due on Friday, March 10.

Instructor: Prof. Artem Zvavitch.

In this HW, we will talk about Lebesgue measure

**Problem 1.** Suppose  $L$  is a linear transformation of  $\mathbb{R}^n$ . Show that if  $E$  is (Lebesgue) measurable subset of  $\mathbb{R}^n$ , then  $L(E)$  is also measurable. You may follow the following steps:

- Note that if  $E$  is compact, so is  $L(E)$ . Hence if  $E$  is  $\mathbf{F}_\sigma$  set (countable union of closed sets), so is  $L(E)$ .
- Show that  $L$  will satisfy

$$|L(x) - L(x')| \leq M|x - x'|$$

for some  $M > 0$  and all  $x, x' \in \mathbb{R}^n$ . Thus  $L$  maps any cube of side length  $l$  into a cube of side length  $2\sqrt{n}Ml$ . Now if  $m(E) = 0$ , there is a collection of cubes  $\{Q_j\}$  such that  $E \subset \cup Q_j$ , and  $\sum m(Q_j) < \varepsilon$ . Thus  $m_*(L(E)) \leq c'\varepsilon$  and hence  $m(L(E)) = 0$ . Combine this with the property that  $E$  is measurable if differs from a  $\mathbf{F}_\sigma$  set by a measure zero.

**Problem 2.** Suppose  $L$  is a linear transformation of  $\mathbb{R}^n$ . Show that if  $E$  is (Lebesgue) measurable subset of  $\mathbb{R}^n$ , then  $m(L(E)) = |\det L|m(E)$

For the next few problems, let  $\Phi = (\Phi_1(x), \Phi_2(x), \dots, \Phi_n(x))$  be a bijection of open set  $O \subset \mathbb{R}^n$  onto another open set  $O' \subset \mathbb{R}^n$ , such that  $\Phi$  is of the class  $C^1$  (i.e. each  $\frac{\partial \Phi_k}{\partial x_i}$ ,  $k, i = 1, 2, \dots, n$  exist and continuous on  $O$ ).

**Problem 3.** Follow ideas proposes for a solution of Problem 1 to show that for every measurable  $E \subset O$  we get  $\Phi(O)$  is measurable.

**Problem 4.** Prove that if  $Q \subset O$  is a cube with diameter  $\varepsilon$  (i.e. the length of the diagonal of the cube is less then  $\varepsilon$ ) and  $z$  is a center of  $Q$  then

$$\Phi(x) = \Phi(z) + \Phi'(z)(x - z) + o(\varepsilon)$$

for all  $x \in Q$ . Note that, using this property we get

$$\Phi(Q) = \Phi(z) + \Phi'(z)(Q - z) + o(\varepsilon).$$

**Problem 5.** Prove that  $m(\Phi(E)) = \int_E |\det \Phi'(x)| dx$ , where  $\Phi'$  is the Jacobian of  $\Phi$ . Here you may assume that  $E$  is a bounded open set, and write  $E = \cup Q_k$ , where  $Q_k$  are cubes whose interiors are disjoint, and whose diameter is less than  $\varepsilon$ . Let  $z_k$  be the center of each  $Q_k$ . Then use the previous problem to show that there exists function  $\eta$  such that  $\eta(\varepsilon) \rightarrow 0$  as  $\varepsilon \rightarrow 0$  such that

$$(1 - \eta(\varepsilon))\Phi'(z_k)(Q_k - z_k) \subset \Phi(Q_k) - \Phi(z_k) \subset (1 + \eta(\varepsilon))\Phi'(z_k)(Q_k - z_k).$$

This would give you

$$m(\Phi(O)) = \sum m(\Phi(Q_k)) = \sum |\det(\Phi'(z_k))| m(Q_k) + o(1), \text{ as } \varepsilon \rightarrow 0,$$

where you, need to use Problem 2 for the second equality.

**Problem 6.** Prove that

$$\int_{O'} f(y) dy = \int_O f(\Phi(x)) |\det \Phi'(x)| dx,$$

for every integrable function  $f$  on  $O'$ .